



HAL
open science

Nitrites and nitrates dietary exposure from natural sources and additives and type-2 diabetes risk

Bernard Srour, Eloi Chazelas, C Debras, N Druesne-Pecollo, C Agaesse, F Szabo de Edelenyi, L Sellem, Emmanuelle Kesse-Guyot, Mélanie Deschasaux-Tanguy, Mathilde Touvier

► To cite this version:

Bernard Srour, Eloi Chazelas, C Debras, N Druesne-Pecollo, C Agaesse, et al.. Nitrites and nitrates dietary exposure from natural sources and additives and type-2 diabetes risk. 15th European Public Health Conference 2022 - Berlin, Germany Strengthening health systems: improving population health and being prepared for the unexpected, Nov 2022, Berlin, Germany. 10.1093/eurpub/ckac129.490 . hal-04071724

HAL Id: hal-04071724

<https://cnam.hal.science/hal-04071724>

Submitted on 17 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

8.D. Oral presentations: Diabetes and nutrition

Abstract citation ID: ckac129.490

Nitrites and nitrates dietary exposure from natural sources and additives and type-2 diabetes risk

Bernard Srour

B Srour¹, E Chazelas¹, C Debras¹, N Druet-Pecollo¹, C Agaesse¹, F Szabo de Edelenyi¹, L Sellem¹, E Kesse-Guyot¹, M Deschasaux-Tanguy¹, M Touvier¹

¹EREN, Inserm, Inrae, Cnam, USPN, INSERM, Bobigny, France
Contact: b.srou@eren.smbh.univ-paris13.fr

Nitrates and nitrites occur naturally in water and soil and are commonly ingested from drinking water and dietary sources. They are also used as food additives. The epidemiological evidence linking exposure to nitrites/nitrates with type-2 diabetes (T2D) risk is scarce. We aimed to study these associations in a large population based prospective cohort study. Overall, 104,168 adults from the French NutriNet-Santé cohort study (median follow-up time 6.7 years) were included. Associations between intakes of nitrites and nitrates (evaluated using repeated 24h dietary records, linked to a comprehensive food composition database and accounting for details of commercial names/brands of industrial products) and risk of T2D were assessed using cause-specific multivariable Cox proportional hazard models adjusted for known risk factors (sociodemographic, anthropometric, lifestyle, medical history,

and nutritional factors). During follow-up, 969 incident T2D cases were ascertained. Total nitrites and nitrates from natural sources were both positively associated with higher T2D risk (HRtertile 3 vs.1 = 1.29 (95% CI 1.06-1.56), Ptrend = 0.004, and 1.27 (95% CI 1.05-1.54), Ptrend = 0.01, respectively). Participants with higher exposure to nitrites from food additives (i.e. above the sex-specific median), and specifically those having higher exposure to sodium nitrite (e250) had a higher T2D risk compared with those who were not exposed to food additive nitrites (HRtertile 3 vs.1 = 1.58 (95% CI 1.28-1.94), Ptrend < 0.001, and 1.59 (95% CI 1.30-1.96), Ptrend < 0.001, respectively). There was no evidence for an association between nitrates of any source and T2D risk (all Ptrend > 0.4). In this large prospective cohort, a higher dietary exposure to nitrites (from both natural sources and food additives) was associated with higher T2D risk. These results provide additional evidence in the context of current discussions about updating regulations on the use of nitrites as food additives.

Key messages:

- A high exposure to dietary nitrites (from both natural and food additive sources) is associated with an increased risk of type-2 diabetes.
- These findings support further regulations concerning the use of nitrites as food additives in processed meats.