



HAL
open science

Adverse Maternal and Infant Outcomes in Women With Chronic Hypertension in France (2010–2018): The Nationwide CONCEPTION Study

Grégory Lailler, Clémence Grave, Amélie Gabet, Nolwenn Regnault, Catherine Deneux-tharoux, Sandrine Kretz, Claire Mounier-vehier, Vassilis Tsatsaris, Geneviève Plu-bureau, Jacques Blacher, et al.

► To cite this version:

Grégory Lailler, Clémence Grave, Amélie Gabet, Nolwenn Regnault, Catherine Deneux-tharoux, et al.. Adverse Maternal and Infant Outcomes in Women With Chronic Hypertension in France (2010–2018): The Nationwide CONCEPTION Study. *Journal of the American Heart Association*, 2023, 12 (5), 10.1161/JAHA.122.027266 . hal-04086582

HAL Id: hal-04086582

<https://cnam.hal.science/hal-04086582>

Submitted on 2 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.



L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

ORIGINAL RESEARCH

Adverse Maternal and Infant Outcomes in Women With Chronic Hypertension in France (2010–2018): The Nationwide CONCEPTION Study

Grégory Lailier , MD; Clémence Grave , MD; Amélie Gabet , PhD; Nolwenn Regnault, PhD; Catherine Deneux-Tharoux , MD, PhD; Sandrine Kretz , MD, PhD; Claire Mounier-Vehier , MD, PhD; Vassilis Tsatsaris , MD, PhD; Geneviève Plu-Bureau , MD, PhD; Jacques Blacher , MD, PhD; Valérie Olié , PhD

BACKGROUND: It has been suggested that chronic hypertension is a risk factor for negative maternal and fetal outcomes during pregnancy and postpartum. We aimed to estimate the association of chronic hypertension on adverse maternal and infant outcomes and assess the impact of antihypertensive treatment and these outcomes.

METHODS AND RESULTS: Using data from the French national health data system, we identified and included in the CONCEPTION cohort all women in France who delivered their first child between 2010 and 2018. Chronic hypertension before pregnancy was identified through antihypertensive medication purchases and by diagnosis during hospitalization. We assessed the incidence risk ratios (IRRs) of maternofetal outcomes using Poisson models. A total of 2822616 women were included, and 42349 (1.5%) had chronic hypertension and 22816 were treated during pregnancy. In Poisson models, the adjusted IRR (95% CI) of maternofetal outcomes for women with hypertension were as follows: 1.76 (1.54–2.01) for infant death, 1.73 (1.60–1.87) for small gestational age, 2.14 (1.89–2.43) for preterm birth, 4.58 (4.41–4.75) for preeclampsia, 1.33 (1.27–1.39) for cesarean delivery, 1.84 (1.47–2.31) for venous thromboembolism, 2.62 (1.71–4.01) for stroke or acute coronary syndrome, and 3.54 (2.11–5.93) for maternal death postpartum. In women with chronic hypertension, being treated with an antihypertensive drug during pregnancy was associated with a significantly lower risk of obstetric hemorrhage, stroke, and acute coronary syndrome during pregnancy and postpartum.

CONCLUSIONS: Chronic hypertension is a major risk factor of infant and maternal negative outcomes. In women with chronic hypertension, the risk of pregnancy and postpartum cardiovascular events may be decreased by antihypertensive treatment during pregnancy.

Key Words: antihypertensive agents ■ blood pressure ■ epidemiology ■ hypertension ■ pregnancy complication

Chronic hypertension in pregnancy, defined as hypertension predating pregnancy or diagnosed before 20 weeks of gestation, concerns 1% to 4% of pregnancies in the United States and 1.7% of pregnancies in France.^{1–3} Previous studies reported an increasing temporal trend in the prevalence of chronic hypertension and other hypertensive disorders of pregnancy, presumably correlated with older maternal age at first birth and

an increased prevalence of obesity.^{1,4–6} It has been suggested that chronic hypertension is a risk factor for negative maternal and fetal outcomes during pregnancy and postpartum^{7–12} and that these correlations are not entirely mediated by the excess risk of preeclampsia.⁸ A recent systematic review and meta-analysis of 94 studies found that chronic hypertension was significantly associated with many negative maternal and perinatal outcomes,

Correspondence to: Grégory Lailier, Santé publique France, Saint-Maurice, France. Email: gregory.lailier@santepubliquefrance.fr

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.122.027266>

For Sources of Funding and Disclosures, see page 9.

© 2023 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- Of primiparous women, 1.5% had chronic hypertension, and 54% of women with hypertension were treated with antihypertensive drugs during pregnancy.
- In primiparous women, chronic hypertension was associated with infant death, small gestational age, preterm birth, preeclampsia, cesarean delivery, venous thromboembolism, stroke or acute coronary syndrome, and maternal death postpartum.
- In women with chronic hypertension, being treated with an antihypertensive drug during pregnancy was associated with a significantly lower risk of obstetric hemorrhage, stroke, and acute coronary during pregnancy and postpartum.

What Are the Clinical Implications?

- Chronic hypertension is a major risk factor of infant and maternal negative outcomes.
- In women with chronic hypertension, the risk of pregnancy and postpartum cardiovascular events may be decreased by antihypertensive treatment during pregnancy.

Nonstandard Abbreviations and Acronyms

CONCEPTION	Cohort of Cardiovascular Diseases in Pregnancy
IRR	incidence risk ratio

including preeclampsia, cesarean delivery, maternal mortality, preterm birth, stillbirth, and small-for-gestational age (SGA).⁷ However, few studies to date have assessed the association between chronic hypertension and maternal death or major cardiovascular events, such as acute coronary syndrome (ACS),^{13,14} stroke,¹⁵ and thromboembolism,¹⁶ during pregnancy and postpartum.

Current guidelines for the management of hypertension and cardiovascular diseases during pregnancy recommend initiating pharmacological treatment in all women with severe hypertension and those with mild to moderate hypertension and a high cardiovascular risk.^{17–19} However, the impact of antihypertensive treatment during pregnancy on maternal and fetal outcomes is debated, and evidence remains scarce.²⁰ The most recent Cochrane review on this topic concluded that antihypertensive drug therapy for mild to moderate hypertension during pregnancy probably halves the risk of developing severe hypertension but may have little

or no effect on other clinically important outcomes, including child and mother death, preeclampsia, preterm birth, and fetal growth restriction.²¹ In the CHAP (Chronic Hypertension and Pregnancy) randomized controlled trial, Tita et al²² found that a strategy of targeting a blood pressure of <140/90 mmHg was associated with better pregnancy outcomes than a strategy of reserving treatment only for severe hypertension, with no increase in the risk of SGA birth weight.

To the best of our knowledge, the impact of antihypertensive treatment for chronic hypertension on the occurrence of major cardiovascular events during pregnancy (ACS, stroke, venous thromboembolism) has never been assessed.

In this context, we aimed to estimate the impact of chronic hypertension on adverse maternal and infant outcomes, including cardiovascular events, in women in France and assess the impact of antihypertensive treatment on these outcomes.

METHODS

Data Source

CONCEPTION (Cohort of Cardiovascular Risk in Pregnancy) is an ongoing prospective cohort including all women resident in France who gave birth in France between January 1, 2010, and December 31, 2018. A detailed description of the cohort protocol is available in previous articles.^{2,23} Cohort data were extracted from the French national health insurance information system database (Système National des Données de Santé),^{21,23,24} which contains comprehensive information on all health care expenditures reimbursed by France's national health insurance system for the entire population. Specifically, it contains information about all public and private hospital stays, including diagnosis and medical interventions, as well as information on outpatient care, including all reimbursements for drug purchases.

In line with the French national regulations and ethics committee, participant consent and institutional review board approval were not required for this study. Santé Publique France—the French public health agency—has full and permanent access to the Système National des Données de Santé (governmental deliberation No. 2016–316, October 13, 2016). We cannot share national health data system data as they are only available on a secure portal. Authorization to access this portal needs registration and clearance.

Study Population

The present study included all first deliveries of women enrolled in the CONCEPTION cohort (hereafter referred to as primiparous) who gave birth in a hospital after 22 weeks of gestation between January 1, 2010, and December 31, 2018. Before 22 weeks of

gestation, fetal losses are defined as miscarriages and are therefore not always identifiable in the *Système National des Données de Santé*. Women who underwent a termination of pregnancy for a maternal or fetal reason were excluded as were those with a history of stroke, ACS, or heart failure after January 1, 2006. The study population was divided into 2 subpopulations according to the presence or not of chronic hypertension before pregnancy. The latter was defined as the dispensing of antihypertensive medication on at least 3 different dates between 1 year preceding the pregnancy and 20 weeks of gestation or on 2 different dates if at least 1 large package of antihypertensive drugs was dispensed or if they were hospitalized with a primary diagnosis of preexisting chronic hypertension (*International Classification of Diseases, Tenth Revision [ICD-10] codes: O10, O11*) during pregnancy or postpartum. Chronic hypertension was considered treated during pregnancy if antihypertensive medication had been dispensed at least once between 20 weeks of gestation and delivery. Antihypertensive treatment initiated after ACS, stroke, or heart failure during pregnancy was presumed to have been prescribed as a secondary prevention treatment and was therefore not considered in the identification of treated hypertension during pregnancy.²

Outcomes

We searched for adverse maternal outcomes from the date of pregnancy to the sixth week postpartum by identifying hospitalizations with the following *ICD-10* codes: delivery and postpartum hemorrhage, hereafter referred to as obstetric hemorrhage (O46, O67, O72), stroke (I60 to I64), venous thromboembolism (I80 to I82, O223, O871, O879, I676, I636, O873, O225, O229, O882, I26), ACS (I20 to I23), and preeclampsia (O14). Gestational diabetes was identified using an algorithm combining the delivery of insulin and glucose strips or a diagnosis of diabetes during pregnancy (E10 to E14, O24) with no preexisting diabetes. Because the study included only women who delivered, maternal death was recorded only during the postpartum period.

The following infant and maternal outcomes and covariates were identified from hospital maternity discharge summaries: multiple pregnancy, delivery mode (vaginal or cesarean delivery), infant death, premature birth defined as a live birth before 37 weeks of gestation and SGA (ie, <10th percentile for sex and gestational age).

Women's Characteristics and Covariates

Women who benefited from Universal Medical Coverage (*Couverture médicale universelle complémentaire*), a social benefit in France for those whose income is below a certain ceiling, were defined as

living in social deprivation. Smoking was identified by specific coding during hospitalization (F17, Z716, Z720, T652), by the reimbursement of payments for nicotine replacement treatments before or during pregnancy, or by a "newborn affected by maternal use of tobacco" (P042) diagnosis. Previous diabetes was identified by the dispensing of at least 3 antidiabetic prescriptions on 3 different dates (or on 2 dates if at least 1 large package of antidiabetic drugs was dispensed) in the year preceding pregnancy. A personal history of venous thromboembolic disease was identified between January 1, 2006, and the date of pregnancy using the criteria cited previously (see Outcomes). Obesity was identified from maternity hospital discharge reports. Gestational age was expressed in completed weeks of amenorrhea.

For women with chronic hypertension before pregnancy, we identified the dispensing of antihypertensive medications, which we divided into the following 5 categories: (1) angiotensin-converting enzyme inhibitors and angiotensin II receptor antagonists, (2) β -blockers, (3) diuretics, (4) calcium channel blockers, and (5) other.

Antihypertensive Drugs During Pregnancy

For women with chronic hypertension treated during pregnancy, we identified the dispensing of antihypertensive drugs at each trimester of pregnancy and during the 6 weeks postpartum period. Angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, and aliskiren were considered contraindicated drugs during pregnancy (formally contraindicated in the second and third trimesters). Methyldopa, nicardipine, labetalol, and nifedipine were considered indicated drugs during pregnancy. All other molecules were considered nonindicated during pregnancy.²⁵

Statistical Analysis

We described mothers' and children's characteristics in the total population and according to a diagnosis of chronic hypertension before pregnancy by calculating numbers and percentages for categorical variables and mean and standard deviation for the age. Maternal and infant outcomes were described according to the existence of chronic hypertension and the treatment of hypertension during pregnancy. The frequency of these outcomes was compared using a χ^2 test between the chronic hypertension and no hypertension groups and between the treated hypertension and untreated hypertension groups. The distribution of different gestational ages at birth was described by curves according to the existence of chronic hypertension and the treatment of hypertension during pregnancy.

We performed univariate and multivariate Poisson regression models to estimate the incidence risk ratios (IRRs) of maternal and infant outcomes according to

chronic hypertension versus no hypertension, treated hypertension versus no hypertension, and untreated hypertension versus no hypertension, with a 95% CI. Multivariate Poisson models were performed corrected for overdispersion and adjusted for maternal age (years), social deprivation, obesity, tobacco use, history of diabetes, multiple pregnancy, gestational diabetes, and preeclampsia. Models estimating the IRRs of gestational diabetes and preeclampsia were adjusted on these covariates except for gestational diabetes.

As a sensibility analysis, we computed a propensity score of being treated during pregnancy for women with chronic hypertension before pregnancy based on a logistic regression model. All previously cited covariates were included in the propensity score plus the year of childbirth. We then estimated the odds ratios of maternal and infant outcomes using logistic regression models (crude and weighted on the inverse probability of treatment). The purpose of this sensibility analysis was to assess whether the associations between mother and child outcomes and antihypertensive treatment during pregnancy persist when a propensity score instead of a multiple adjustment controls the indication bias.

For women with chronic hypertension treated during pregnancy, we calculated the numbers and percentages of women who took only indicated antihypertensive drugs, nonindicated drugs, or contraindicated drugs.

RESULTS

We identified 2 833 376 primiparous women who gave birth in France during the period from January 1, 2010, to December 31, 2018 (Figure 1). After the exclusion of 7847 terminations of pregnancy for maternal or fetal medical indication and 2913 women with histories of stroke, ACS, or heart failure, the analysis population included 2 822 616 women. Of these, 42 349 (1.5%) had chronic hypertension before pregnancy, and 22 816 were treated for this condition during pregnancy (54% of women with chronic hypertension). Table 1 shows the population's characteristics according to the existence of chronic hypertension before pregnancy. Women with chronic hypertension were, on average, >3 years older and were more likely to smoke, live in social deprivation, be obese and diabetic, and have previous venous thromboembolism medical history.

All pregnancy and maternal outcomes were statistically more frequent in women with chronic hypertension than in women without chronic hypertension (Table 2). Singletons born in 2013 or after had mean birth weights of 3214.5 g and 2963.8 g for mothers without and with chronic hypertension, respectively (mean difference = 251 g).




Figure 1. Study flowchart.

In fully adjusted Poisson models (Figure 2), the IRRs (95% CI) of infant outcomes for women with chronic hypertension were 1.76 (1.54–2.01) for infant death, 1.73 (1.60–1.87) for SGA, and 2.14 (1.89–2.43) for preterm birth compared with women without chronic hypertension. The adjusted IRRs for maternal outcomes were 1.29 (1.25–1.33) for gestational diabetes, 4.58 (4.41–4.75) for preeclampsia, 1.33 (1.27–1.39) for cesarean delivery, 1.09 (1.05–1.14) for obstetric hemorrhage, 1.84 (1.47–2.31) for venous thromboembolism, 2.67 (1.71–4.01) for stroke or ACS, and 3.54 (2.11–5.93) for maternal death postpartum. IRRs were higher in women with treated hypertension during pregnancy for gestational diabetes and preeclampsia. Figure 3 shows the frequency curves of gestational age at birth according to the existence of chronic hypertension and hypertension treatment during pregnancy. Women with chronic hypertension gave birth on average 1 week earlier than women without chronic hypertension (37.9 weeks of amenorrhea versus 39.0 weeks).

In women with chronic hypertension, being treated with an antihypertensive drug during pregnancy was associated with a significant excess risk (adjusted IRR [95% CI]) of SGA (1.47 [1.39–1.57]), preterm birth (1.26 [1.17–1.36]), cesarean delivery (1.17 [1.14–1.20]), gestational diabetes (1.25 [1.21–1.30]), and preeclampsia (1.08 [1.04–1.13]) but not of infant death (1.10 [0.87–1.38]). On the contrary, being treated with an antihypertensive drug during pregnancy was associated with a decreased risk of hemorrhage, venous thromboembolism, and stroke or ACS during pregnancy and postpartum. However, this difference only reached

Table 1. Population Characteristics and Dispensing of Antihypertensive Drugs

	Total, N=2822616		Chronic hypertension before pregnancy			
			No, n=2780267		Yes, n=42349	
	No. or mean	Percentage or SD	No. or mean	Percentage or SD	No. or mean	Percentage or SD
Maternal characteristic						
Maternal age, y	28.29	(5.38)	28.29	(5.36)	28.29	(6.01)
Multiple pregnancy	56544	2.00	55210	1.99	1334	3.15
Smoking	251 303	8.90	246635	8.87	4668	11.02
Social deprivation	389005	13.78	382864	13.77	6141	14.50
Medical history						
Obesity	116556	4.13	110410	3.97	6146	14.51
Diabetes	14510	0.51	12783	0.46	1727	4.08
Previous venous thromboembolism	3491	0.12	3321	0.12	170	0.40
Antihypertensive drugs dispensing before 20 wks of gestation*						
ACE inhibitors/AIIRA	8078	19.07
β-blockers	22 171	52.35
Diuretics	3696	8.73
Calcium channel blockers	8194	19.35
Other	5908	13.95
None	33 463	79.02

SD data are shown in parentheses. ACE indicates angiotensin-converting enzyme; and AIIRA, angiotensin II receptor antagonist.

*Number of women diagnosed with chronic hypertension before pregnancy with at least 1 dispensing of antihypertensive drugs between 1 year before the date of pregnancy and 20 weeks of amenorrhea.

statistical significance for hemorrhage (0.93 [0.86–0.99]) and the composite end point “stroke or ACS” (0.38 [0.18–0.83]). In a sensitivity analysis in which the 2913 women with a medical history of stroke, ACS, or heart failure were not excluded, these lower adjusted IRRs were not significant (data not shown).

In women with chronic hypertension, the probability of being treated during pregnancy was estimated using a logistic regression, which comprised all previously cited covariates plus the year of childbirth. The density of the propensity score showed an important overlap between the treated and untreated groups (Figure S1). The associations between antihypertensive treatment and mother and child outcomes estimated by logistic regression weighted on inverse probability of treatment were similar to those estimated by adjusted Poisson regression models (Table S1).

With respect to the 22816 women treated for hypertension, an antihypertensive drug was dispensed for 13795 (60.5%) of them during the first trimester, 16473 (72.2%) during the second trimester, and 14779 (64.8%) during the third trimester. A contraindicated drug (angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, or aliskiren) was dispensed for 328 (2.0%) of all 22816 women during the second trimester and for 105 (0.7%) women during the third trimester (Table 3).

DISCUSSION

In the large-scale nationwide prospective French cohort CONCEPTION, which included, among others, 2833376 primiparous women who gave birth between 2010 and 2018, we found that chronic hypertension before pregnancy was an important risk factor for adverse pregnancy outcomes, cardiovascular diseases, and death both during pregnancy and postpartum. This excess risk was significant irrespective of whether hypertension was treated during pregnancy. When treated during pregnancy, women with chronic hypertension had a significantly lower risk of obstetric hemorrhage and stroke or ACS than their untreated counterparts.

Although several studies have found an association between chronic hypertension and adverse infant outcomes,^{7–9,11,26,27} evidence concerning the association between chronic hypertension and cardiovascular diseases during pregnancy is scarce.^{7,10} We found that chronic hypertension was associated with preeclampsia, obstetric hemorrhage, thromboembolic venous disease, and stroke/ACS. It was also associated with a 3½ times higher risk of maternal death during postpartum, which is consistent with previous reports.^{6,7,10} Although the absolute risks of major cardiovascular events during pregnancy and postpartum remain low,

Table 2. Numbers and Proportions of Infant and Maternal Adverse Events, According to Hypertension Status

	No hypertension, n=2780267		Chronic hypertension, n=42349		Chronic hypertension treated during pregnancy, n=22816		Chronic hypertension untreated during pregnancy, n=19533		Chronic hypertension vs no hypertension, P value	Treated hypertension vs untreated hypertension, P value
	No.	Percentage	No.	Percentage	No.	Percentage	No.	Percentage		
Infant events										
Infant death	12136	0.44	355	0.84	201	0.88	154	0.79	<0.0001	0.2977
Small-for-gestational age*	220449	13.96	5203	24.38	3223	28.77	1980	19.54	<0.0001	<0.0001
Premature delivery threat	277722	9.99	5514	13.02	3288	14.41	2226	11.40	<0.0001	<0.0001
Preterm birth†	201767	7.30	8010	19.11	4857	21.52	3153	16.30	<0.0001	<0.0001
Preterm birth stages†									<0.0001	<0.0001
Moderate preterm (≥32 WG)	170722	6.17	6093	14.54	3682	16.32	2411	12.47		
Very preterm (27–31 WG)	24271	0.88	1569	3.74	983	4.36	586	3.03		
Extremely preterm (<27 WG)	6774	0.24	348	0.83	192	0.85	156	0.81		
Maternal events										
Gestational diabetes	229593	8.26	7110	16.79	4547	19.93	2563	13.12	<0.0001	<0.0001
Preeclampsia	75222	2.71	7237	17.09	4124	18.08	3113	15.94	<0.0001	<0.0001
Cesarean delivery	736920	26.51	20864	49.27	12610	55.27	8254	42.26	<0.0001	<0.0001
Obstetric hemorrhage	158957	5.72	2982	7.04	1582	6.93	1400	7.17	<0.0001	0.35
Venous thromboembolism	3281	0.12	121	0.29	62	0.27	59	0.30	<0.0001	0.56
ACS or stroke	770	0.03	45	0.11	16	0.07	29	0.15	<0.0001	0.01
Death postpartum	152	0.01	13	0.03	9	0.04	4	0.02	<0.0001	0.27

ACS indicates acute coronary syndrome; and WG, weeks of gestation.

*Birth weight was available from 2013; percentages for small for gestational age are calculated for available data for the period from 2013 to 2018.

†Preterm percentages are calculated for live births.




Figure 2. Adjusted IRRs of infant (A) and maternal (B) outcomes according to prepregnancy hypertension and the treatment of hypertension during pregnancy.

IRRs of preeclampsia and gestational diabetes were adjusted for maternal age (years), deprivation (Couverture médicale universelle complémentaire), obesity, tobacco use, history of diabetes, and multiple pregnancy. Other IRRs were adjusted for factors cited previously plus gestational diabetes. Birth weight was available from 2013, low birth weight IRRs were calculated for available data at the period from 2013 to 2018. Preterm IRRs were calculated for live births. ACS indicates acute coronary syndrome; HT, hypertension; IRR, incidence risk ratio; PP, postpartum; and WG, weeks of gestation.

these events have dramatic consequences on both mothers and children, including physical or mental disabilities and death. They have become the main cause of maternal mortality in developed countries.^{28,29} Given the increasing trend of chronic hypertension before pregnancy, this global public health issue is likely to worsen in the next few years, which highlights the need to improve the prevention, screening, and multidisciplinary management of chronic hypertension in women of childbearing age.

Previous studies reported that antihypertensive treatment is beneficial in preventing several negative maternal outcomes.^{7,21,30} Particularly, the CHIPS trial (Control of Hypertension in Pregnancy Study) found that tight versus less-tight control of hypertension in pregnancy did not improve maternal and infant

outcomes, except the risk of severe maternal hypertension, thrombocytopenia, and elevated liver enzymes.³⁰ However, a post hoc analysis of CHIPS data found that severe hypertension was associated with all outcomes except for maternal readmission. Recently, the CHAP trial found that, in women with chronic hypertension, a pharmacological treatment targeting a blood pressure of <140/90 mmHg was associated with a decreased risk of a composite outcome combining severe preeclampsia, medically indicated preterm birth, placental abruption, or fetal or neonatal death.²²

In our study, antihypertensive treatment during pregnancy was significantly associated with a higher risk of cesarean delivery, SGA, and preterm birth but not of infant death. Nevertheless, it remains unclear whether antihypertensive drugs have a negative causal




Figure 3. Distribution of gestational age at birth according to chronic hypertension and antihypertensive treatment. HT indicates hypertension.

effect on pregnancy outcomes or if they only reflect the severity of hypertension. Indeed, we were not able to adjust our models for the severity of hypertension because blood pressure measurements are not available for the cohort. Among all studied outcomes, the association between antihypertensive treatment and SGA was the strongest and could be partly explained by the prescription of β -blockers, which has already been associated with fetal growth restriction.^{31,32}

Moreover, we found that being treated with an antihypertensive drug during pregnancy was associated with a significantly lower risk of delivery or postpartum hemorrhage and of stroke/ACS during pregnancy and postpartum. To the best of our knowledge, this is the first study to report such results. Nevertheless, they must be considered with caution. Given the observational design of this study, a causal relationship

between the treatment of chronic hypertension during pregnancy and a lower risk of hemorrhage or stroke cannot be asserted. In a sensibility analysis using a propensity score, these associations were unchanged, but the discriminating power of this propensity score was flawed by the lack of blood pressure measurements.

Nonetheless, this finding remains valuable because a clinical trial would require a considerable number of participants to assess the effect of antihypertensive treatment on such rare events and therefore would be unethical and difficult to conduct.

Our results highlight the importance of preconceptional care for women with chronic hypertension to optimize pregnancy planning. This would allow providing women information about the excess risk of negative maternal and infant outcomes and the importance of

Table 3. Number and Type of Antihypertensive Drugs Dispensed to Women With Chronic Hypertension Treated During Pregnancy at All 3 Trimesters of the Pregnancy and Postpartum

Indication*	1 year before pregnancy		First trimester		Second trimester		Third trimester		Postpartum	
	No.	Percentage	No.	Percentage	No.	Percentage	No.	Percentage	No.	Percentage
Contraindicated drug	4563	31.7	1498	10.9	328	2.0	105	0.7	1584	12.8
Nonindicated drug	8684	60.3	7144	51.8	7022	42.6	5013	33.9	4382	35.3
Indicated drug	1143	7.9	5153	37.4	9123	55.4	9661	65.4	6436	51.9
All drugs	14390	100	13795	100	16473	100	14779	100	12402	100

*Angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, and aliskiren were considered contraindicated drugs during pregnancy (formally contraindicated in the second and third trimesters). Methyldopa, nifedipine, labetalol, and nifedipine were considered indicated drugs during pregnancy. All other molecules were considered nonindicated during pregnancy.

blood pressure control during the pregnancy, especially considering the results of the CHIPS and CHAP clinical trials. Our findings also support previous recommendations that women with chronic hypertension be monitored closely for the potential development of adverse complications during pregnancy.

A contraindicated treatment (angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, or aliskiren) had been dispensed during the second or third trimesters of pregnancy for a small proportion of the women in our study. These drugs can lead to fetal complications or death. The importance of stopping or substituting these drugs for others during pregnancy should therefore be emphasized for both mothers and clinicians.¹⁹

This study has many strengths. The use of a national database enabled us to create a near-exhaustive nationwide cohort of women who gave birth in France at some point during a 9-year time period and to avoid inclusion bias. The large sample size of this cohort ensured optimal statistical power to study rare events such as maternal cardiovascular events and death during pregnancy or postpartum, compare treated and untreated pregnant women with chronic hypertension, and adjust our models for several covariates to limit confounding bias. The combination of data on hospital-based diagnoses and outpatient drug dispensing enabled us to study not only infant outcomes but also maternal cardiovascular events and analyze the consumption of each antihypertensive drug class during pregnancy and postpartum. Moreover, the use of hospital records and drug-dispensing data to identify outcomes (diagnoses and treatments) lowered the risk of classification bias, such as recall bias.

This study also had limitations. Because chronic hypertension was identified by the dispensing of antihypertensive drugs and hospital diagnoses, untreated hypertension not reported during hospitalization may have been missed and therefore underestimated. Moreover, we cannot exclude the possibility that a small proportion of women may have taken antihypertensive drugs for indications other than hypertension (eg, kidney failure, migraine). Having said that, apart from migraine, such indications are rare in young women. These possible misclassifications would likely lead to an underestimation of the IRRs of perinatal and maternal outcomes in women with chronic hypertension. Although we comprehensively identified every medication dispensing event, we can only assume that these treatments were actually taken. This assumption could therefore lead to an underestimation of the potential effect of antihypertensive drugs on perinatal and maternal outcomes. Likewise, the compliance to antihypertensive drugs could not be assessed because we could not know how many drugs women were taking. Moreover, the *Système National des Données de*

Santé is a medico-administrative database and therefore lacks clinical data such as weight or blood pressure, resulting in residual confounding. Similarly, no data are available for rates of postpartum hypertension and rates of readmissions in the postpartum period. Finally, as our analysis was conducted on women who delivered for the first time, these results cannot be generalized to subsequent pregnancies.

CONCLUSIONS

Chronic hypertension is a major risk factor of perinatal negative outcomes, including infant death and maternal venous thromboembolic or cardiovascular events during pregnancy and postpartum. Women treated with antihypertensive drugs during pregnancy were at higher risk of negative perinatal outcomes and at lower risk of postpartum cardiovascular events, yet no causal inference can be established between the antihypertensive treatment and these outcomes given the observational design of this study. Further studies should be done on how antihypertensive therapy affects not just pregnancy outcomes but also long-term outcomes such as ACS, cardiovascular diseases, and death among these women.

ARTICLE INFORMATION

Received August 4, 2022; accepted November 1, 2022.

Affiliations

Santé publique France, Saint-Maurice, France (G.L., C.G., A.G., N.R., V.O.); Department of Medicine, Université Paris Est, Créteil, France (G.L.); Obstetrical Perinatal and Pediatric Epidemiology Research Team (EPOPé), Centre for Epidemiology and Statistics Sorbonne Paris Cité (CRESS), French National Institute for Health and Medical Research (INSERM), Paris, France (C.D., G.P.); Department of Medicine, Université Paris Cité, Paris, France (C.D., V.T., G.P., J.B.); Centre de diagnostic et de thérapeutique, Assistance Publique Hôpitaux de Paris (AP-HP), Hôtel Dieu de Paris, Paris, France (S.K., J.B.); CHU Lille, Institut Cœur-Poumon, Médecine Vasculaire et HTA, Lille, France (C.M.); Maternité Port-Royal, Assistance Publique Hôpitaux de Paris (AP-HP), Hôpital Cochin, Paris, France (V.T.); and Unité de gynécologie médicale, Assistance Publique Hôpitaux de Paris (AP-HP), Hôpital Port-Royal Cochin, Paris, France (G.P.).

Sources of Funding

This work was supported by the French Hypertension Society, the Hypertension Research Foundation, and the French Cardiology Federation through the call for scientific projects titled "Thematic Grant 2019: Cardiovascular Diseases in Women." The funders had no role in the study design, data collection, data analysis, decision to publish, or drafting of the manuscript.

Disclosures

Dr Kretz reports, outside the submitted work, nonfinancial support from Lilly France, Novonordisk, Novartis Pharma, Roche diabetes care, Lifescan, Abbott France, Sanofi, ViiV Healthcare, Servier, and Becton Dickinson and personal fees from Icomed, Pascaleo, BT3SI, and M3global research. Dr Mounier-Vehier reports, outside the submitted work, personal fees and nonfinancial support from Servier, Lundbeck, Astrazeneca, Merck & Co, and Boehringer. Dr Blacher reports, outside the submitted work, personal fees from Abbott, Bayer, Böttu, Ferring, Steripharma, Kantar, and Teriak; personal fees and nonfinancial support from Pfizer and Quantum Genomics; and personal fees from Sanofi and Servier. All other authors declare no conflict of interest.

Supplemental Material

Table S1
Figure S1

REFERENCES

- Hutcheon JA, Lisonkova S, Joseph KS. Epidemiology of pre-eclampsia and the other hypertensive disorders of pregnancy. *Best Pract Res Clin Obstet Gynaecol*. 2011;25:391–403. doi: 10.1016/j.bpobgyn.2011.01.006
- Olie V, Moutengou E, Grave C, Deneux-Tharoux C, Regnault N, Kretz S, Gabet A, Mounier-Vehier C, Tsatsaris V, Plu-Bureau G, et al. Prevalence of hypertensive disorders during pregnancy in France (2010–2018): the Nationwide CONCEPTION Study. *J Clin Hypertens (Greenwich)*. 2021;23:1344–1353. doi: 10.1111/jch.14254
- Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, Hall DR, Warren CE, Adoyi G, Ishaku S, et al. Hypertensive disorders of pregnancy: ISSHP classification, diagnosis, and management recommendations for international practice. *Hypertension*. 2018;72:24–43. doi: 10.1161/HYPERTENSIONAHA.117.10803
- Blondel B, Lelong N, Kermarrec M, Goffinet F; National Coordination Group of the National Perinatal S. Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *J Gynecol Obstet Biol Reprod*. 2012;41:e1–e15. doi: 10.1016/j.jgyn.2012.04.014
- Kuklina EV, Ayala C, Callaghan WM. Hypertensive disorders and severe obstetric morbidity in the United States. *Obstet Gynecol*. 2009;113:1299–1306. doi: 10.1097/AOG.0b013e3181a45b25
- Bateman BT, Bansil P, Hernandez-Diaz S, Mhyre JM, Callaghan WM, Kuklina EV. Prevalence, trends, and outcomes of chronic hypertension: a nationwide sample of delivery admissions. *Am J Obstet Gynecol*. 2012;206(134):e131–e138. doi: 10.1016/j.ajog.2011.10.878
- Al Khalaf SY, O'Reilly EJ, Barrett PM, BL DF, Pawley LC, FP MC, Khashan AS. Impact of chronic hypertension and antihypertensive treatment on adverse perinatal outcomes: systematic review and meta-analysis. *J Am Heart Assoc*. 2021;10:e018494. doi: 10.1161/JAHA.120.018494
- McCowan LM, Buist RG, North RA, Gamble G. Perinatal morbidity in chronic hypertension. *Br J Obstet Gynaecol*. 1996;103:123–129. doi: 10.1111/j.1471-0528.1996.tb09662.x
- Bramham K, Parnell B, Nelson-Piercy C, Seed PT, Poston L, Chappell LC. Chronic hypertension and pregnancy outcomes: systematic review and meta-analysis. *BMJ*. 2014;348:g2301. doi: 10.1136/bmj.g2301
- Gilbert WM, Young AL, Danielsen B. Pregnancy outcomes in women with chronic hypertension: a population-based study. *J Reprod Med*. 2007;52:1046–1051.
- Rey E, Couturier A. The prognosis of pregnancy in women with chronic hypertension. *Am J Obstet Gynecol*. 1994;171:410–416. doi: 10.1016/0002-9378(94)90276-3
- Li F, Wang T, Chen L, Zhang S, Chen L, Qin J. Adverse pregnancy outcomes among mothers with hypertensive disorders in pregnancy: a meta-analysis of cohort studies. *Pregnancy Hypertens*. 2021;24:107–117. doi: 10.1016/j.preghy.2021.03.001
- Ladner HE, Danielsen B, Gilbert WM. Acute myocardial infarction in pregnancy and the puerperium: a population-based study. *Obstet Gynecol*. 2005;105:480–484. doi: 10.1097/01.AOG.0000151998.50852.31
- James AH, Jamison MG, Biswas MS, Branciazio LR, Swamy GK, Myers ER. Acute myocardial infarction in pregnancy: a United States population-based study. *Circulation*. 2006;113:1564–1571. doi: 10.1161/CIRCULATIONAHA.105.576751
- Karjalainen L, Tikkanen M, Rantanen K, Aarnio K, Korhonen A, Saaros A, Laivuori H, Gissler M, Ijas P. Stroke in pregnancy and puerperium: validated incidence trends with risk factor analysis in Finland 1987–2016. *Neurology*. 2021;96:e2564–e2575. doi: 10.1212/WNL.0000000000011990
- James AH, Jamison MG, Branciazio LR, Myers ER. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *Am J Obstet Gynecol*. 2006;194:1311–1315. doi: 10.1016/j.ajog.2005.11.008
- Regitz-Zagrosek V, Roos-Hesselink JW, Bauersachs J, Blomstrom-Lundqvist C, Cifkova R, De Bonis M, Iung B, Johnson MR, Kintscher U, Kranke P, et al. 2018 ESC guidelines for the management of cardiovascular diseases during pregnancy. *Eur Heart J*. 2018;39:3165–3241. doi: 10.1093/eurheartj/ehy340
- Mounier-Vehier C, Amar J, Boivin J-M, Denolle T, Fauvel J-P, Plu-Bureau G, Tsatsaris V, Blacher J. Hypertension and pregnancy. Expert consensus statement from the French Society of Hypertension, an affiliate of the French Society of Cardiology. *Presse Med*. 2016;45:682–699. doi: 10.1016/j.jpm.2016.05.012
- American College of O, Gynecologists' Committee on Practice B-O. ACOG practice bulletin No. 203: chronic hypertension in pregnancy. *Obstet Gynecol*. 2019;133:e26–e50. doi: 10.1097/AOG.0000000000003020
- Garovic VD, Dechend R, Easterling T, Karumanchi SA, McMurtry Baird S, Magee LA, Rana S, Vermunt JV, August P; American Heart Association Council on H, et al. Hypertension in pregnancy: diagnosis, blood pressure goals, and pharmacotherapy: a scientific statement from the American Heart Association. *Hypertension*. 2022;79:e21–e41. doi: 10.1161/HYP.0000000000000208
- Abalos E, Duley L, Steyn DW, Gialdini C. Antihypertensive drug therapy for mild to moderate hypertension during pregnancy. *Cochrane Database Syst Rev*. 2018;10:CD002252. doi: 10.1002/14651858.CD002252.pub4
- Tita AT, Szychowski JM, Boggess K, Dugoff L, Sibai B, Lawrence K, Hughes BL, Bell J, Aagaard K, Edwards RK, et al. Treatment for mild chronic hypertension during pregnancy. *N Engl J Med*. 2022;386:1781–1792. doi: 10.1056/NEJMoa2201295
- Boucheron P, Lailler G, Moutengou E, Regnault N, Gabet A, Deneux-Tharoux C, Kretz S, Grave C, Mounier-Vehier C, Tsatsaris V, et al. Hypertensive disorders of pregnancy and onset of chronic hypertension in France: the nationwide CONCEPTION study. *Eur Heart J*. 2021;43:3352–3361. doi: 10.1093/eurheartj/ehab686
- Tuppin P, de Roquefeuil L, Weill A, Ricordeau P, Merliere Y. French national health insurance information system and the permanent beneficiaries sample. *Rev Epidemiol Sante Publique*. 2010;58:286–290. doi: 10.1016/j.respe.2010.04.005
- Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, de Simone G, Dominiczak A, et al. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J*. 2018;39:3021–3104. doi: 10.1093/eurheartj/ehy339
- Fisher SC, Van Zutphen AR, Romitti PA, Browne ML; National Birth Defects Prevention S. Maternal hypertension, antihypertensive medication use, and small for gestational age births in the national birth defects prevention study, 1997–2011. *Matern Child Health J*. 2018;22:237–246. doi: 10.1007/s10995-017-2395-8
- Al Khalaf SY, O'Reilly EJ, McCarthy FP, Kublickas M, Kublickiene K, Khashan AS. Pregnancy outcomes in women with chronic kidney disease and chronic hypertension: a National cohort study. *Am J Obstet Gynecol*. 2021;225:298.e1–298.e20. doi: 10.1016/j.ajog.2021.03.045
- Kotit S, Yacoub M. Cardiovascular adverse events in pregnancy: a global perspective. *Glob Cardiol Sci Pract*. 2021;2021:e202105. doi: 10.21542/gcsp.2021.5
- Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *Lancet*. 2006;367:1066–1074. doi: 10.1016/S0140-6736(06)68397-9
- Magee LA, von Dadelszen P, Rey E, Ross S, Asztalos E, Murphy KE, Menzies J, Sanchez J, Singer J, Gafni A, et al. Less-tight versus tight control of hypertension in pregnancy. *N Engl J Med*. 2015;372:407–417. doi: 10.1056/NEJMoa1404595
- Duan L, Ng A, Chen W, Spencer HT, Lee MS. Beta-blocker subtypes and risk of low birth weight in newborns. *J Clin Hypertens (Greenwich)*. 2018;20:1603–1609. doi: 10.1111/jch.13397
- Kayser A, Beck E, Hoeltzenbein M, Zinke S, Meister R, Weber-Schoendorfer C, Schaefer C. Neonatal effects of intrauterine metoprolol/bisoprolol exposure during the second and third trimester: a cohort study with two comparison groups. *J Hypertens*. 2020;38:354–361. doi: 10.1097/HJH.0000000000002256

SUPPLEMENTAL MATERIAL


Table S1. Unadjusted and weighted Odds Ratios (ORs) of child and mother events in women with chronic hypertension, according the treatment of HT during pregnancy

	Treated vs untreated HT (OR)	
	n=42,329	
	Crude	Weighed on Inverse probability of Treatment*
	OR [95% confidence interval]	OR [95% confidence interval]
Infant events		
Fetal or neonatal death	1.12 [0.91-1.38]	1.13 [0.97-1.31]
Small for gestational age†	1.66 [1.56-1.77]	1.66 [1.59-1.74]
Preterm‡	1.41 [1.34-1.48]	1.33 [1.28-1.38]
Preterm (stages)		
Moderate preterm	1.40 [1.32-1.48]	1.32 [1.27-1.38]
Very preterm	1.53 [1.38-1.70]	1.41 [1.31-1.52]
Extremely preterm	1.13 [0.91-1.39]	1.09 [0.94-1.26]
Maternal events		
Preeclampsia	1.16 [1.11-1.23]	1.11 [1.07-1.15]
Cesarean section	1.69 [1.62-1.76]	1.32 [1.28-1.35]
Delivery or post-partum hemorrhage	0.97 [0.90-1.04]	0.93 [0.88-0.98]
Thromboembolic venous disease	0.90 [0.63-1.29]	0.80 [0.62-1.03]
Stroke or ACS	0.47 [0.26-0.87]	0.37 [0.24-0.58]
Death during post partum	1.93 [0.59-6.26]	1.41 [0.66-3.02]

*The probability of being treated was evaluated by a logistic regression model using the following variates: maternal age (years), deprivation, obesity, tobacco use, history of diabetes, gestational hypertension, history of thromboembolic disease, multiple pregnancy and year of childbirth. †Birth weight was available from 2013, low birth weight ORs are calculated for available data at the period 2013-2018. ‡Preterm ORs are calculated for live births.

Abbreviations: ACS = Acute coronary syndrome, HT = chronic hypertension before pregnancy, OR = Odds Ratio.

Figure S1. Propensity score distribution by treatment groups



The probability of being treated was evaluated by a logistic regression model using the following variates: maternal age (years), deprivation, obesity, tobacco use, history of diabetes, gestational hypertension, history of thromboembolic disease, multiple pregnancy and year of childbirth.